

CMB-NORMALIZED PREDICTIONS FOR SUNYAEV-ZEL'DOVICH EFFECT FLUCTUATIONS

GILBERT P. HOLDER

School of Natural Sciences, Institute for Advanced Study, Princeton NJ, 08540; holder@ias.edu

Submitted to ApJL July 18, 2002

ABSTRACT

We predict the level of small-scale anisotropy in the cosmic microwave background (CMB) due to the Sunyaev–Zel’dovich (SZ) effect for the ensemble of cosmological models that are consistent with current measurements of large-scale CMB anisotropy. We argue that the recently reported detections of the small-scale (arcminutes) CMB anisotropy are only marginally consistent with being the SZ effect when cosmological models are calibrated to the existing primary CMB data on large scales. The discrepancy is at more than $2-2.5\sigma$, and is mainly due to a lower $\sigma_8 \lesssim 0.8$ favored by the primary CMB and a higher $\sigma_8 \gtrsim 1$ favored by the SZ effect. A degeneracy between the optical depth to Thomson scattering and the CMB-derived value of σ_8 suggests that the discrepancy is reduced if the universe was reionized very early, at redshift of ~ 25 .

Subject headings: cosmic microwave background — cosmological parameters — cosmology: observations

1. INTRODUCTION

The current generation of cosmic microwave background (CMB) experiments are producing a wealth of information on a wide range of angular scales. With the recent measurements by BIMA (Dawson et al. 2002) and CBI (Mason et al. 2002) complementing earlier measurements (see Hu and Dodelson 2002, for a recent compilation), CMB anisotropy has been measured over a range of multipoles of $l = 2 - 6000$ (or angular scales of $2' - 90^\circ$).

At low multipoles ($l \lesssim 2000$) anisotropy is primarily generated at $z \gtrsim 1000$ except at very low multipoles ($l \lesssim 10$) where late-time decay of gravitational potential contributes significantly. At higher multipoles (smaller angular scales) low-redshift sources generate a significant amount of fluctuation power. At the observing frequencies of CBI and BIMA (~ 30 GHz), the largest sources of low-redshift anisotropy are radio point sources and the thermal Sunyaev–Zel’dovich (SZ) effect. The latter is of cosmological interest and may be large enough to be detected, depending on cosmology (e.g., Cole and Kaiser 1988).

The reported detections of power at small angular scales ($l = 2000 - 6000$) have argued that point-source contamination is not a problem, suggesting that the detected power could be due to the SZ effect (Bond et al. 2002; Dawson et al. 2002; Komatsu and Seljak 2002). Since the number density and brightness of the sources contributing to the SZ fluctuations (i.e., hot gas in galaxy clusters at $z \lesssim 1$) depend on the background cosmology, the level of the SZ fluctuations depends on cosmological parameters. The SZ angular power spectrum is sensitive to the matter-fluctuation amplitude and the baryon density of the universe but relatively insensitive to the matter density of the universe (Komatsu and Kitayama 1999) or other cosmological parameters (Komatsu and Seljak 2002). By fitting the CBI and BIMA data to theoretical predictions, Komatsu and Seljak (2002) have found a constraint on linear r.m.s. mass fluctuations within an $8 h^{-1}$ Mpc sphere, σ_8 , as $\sigma_8(\Omega_b h/0.035)^{0.29} = 1.04 \pm 0.12$ at the 95% confidence level.

On the other hand, observations of the primary CMB anisotropy and the large-scale structure (LSS) of the universe have already provided tight constraints on cosmological parameters (e.g., Wang et al. 2002). By using these constraints we

can predict how much SZ power ought to be seen at the CBI and BIMA multipole bands. By doing so we can see if an SZ interpretation of the small-scale fluctuations is consistent with cosmological models favored by CMB or LSS.

In this *Letter* we estimate the level of the SZ angular power spectrum expected from cosmological models consistent with CMB data at $l < 2000$, and compare it with the CBI and BIMA data at $2000 < l < 10000$. We use only CMB data and a prior on the Hubble constant and do not include any constraints from LSS. For the primary CMB data, we use the Markov Chain Monte Carlo (MCMC) results of Lewis and Bridle (2002) as an estimate of the appropriate weighting for each cosmological model. For the SZ effect, we use a model for the SZ power spectrum as a function of cosmology from Komatsu and Seljak (2002). In §2 we outline salient features of the MCMC results. In §3 we sketch out our calculation of the SZ power spectrum, including the effects of the non-Gaussian nature of the SZ fluctuations. In §4 we compare MCMC realizations of the SZ power spectrum weighted by the primary CMB data with the CBI and BIMA data, and check for consistency between them. In §5 we discuss implications for the reported detections of power at small angular scales.

2. CMB-CALIBRATED COSMOLOGICAL MODELS

As a subset of cosmological models consistent with the current CMB data, we use the MCMC results of Lewis and Bridle (2002) (for details see <http://cosmologist.info/cosmomc>). Estimating cosmological parameters using MCMC methods (Christensen et al. 2001) entails taking random steps in parameter space, accepting a step if the new point is more likely, otherwise accepting the step with some probability (less than one) set by how much worse the new point is than the current point. The resulting list of points should sample the likelihood function in the multidimensional parameter space.

An advantage of MCMC methods (besides often being faster than grids) is that the resulting list of consistent models (the “chain”) provides a self-consistent sampling of the likelihood distribution. A distribution of CMB-derived cosmological parameters using the MCMC chain therefore provides a maximally informative CMB prior. Covariances between parameters are naturally incorporated. The chain provides more samples

where the density is higher; this feature is actually a disadvantage when dealing with models lying in the tail of the distribution, as we will return to later.

The chain that we use (Lewis and Bridle 2002) assumes a flat universe, no tensor component, and an equation of state for the dark energy component of $w = -1$ (i.e., assumes a cosmological constant rather than quintessence). A prior on the Hubble constant ($h = 0.72 \pm 0.08$) consistent with the Hubble Key Project results (Freedman et al. 2001) is assumed, but no LSS priors. There are six free parameters in the chain: $\Omega_b h^2$, $\Omega_{\text{CDM}} h^2$, Ω_Λ , z_{re} , n_s , and A_s , where Ω_b is the baryon density relative to the critical density, Ω_{CDM} the cold dark-matter density, Ω_Λ the cosmological-constant energy density, z_{re} the redshift of reionization, n_s the scalar spectral index, and A_s the amplitude of density fluctuations. The constraint that spatial curvature is zero defines the Hubble constant as $h = [(\Omega_b + \Omega_{\text{CDM}})h^2 / (1 - \Omega_\Lambda)]^{1/2}$. The total matter density $\Omega_{\text{CDM}} + \Omega_b$ and σ_8 are derived parameters for each model. There are 2,596 MCMC samples in total.

For each of the cosmological models in the chain, we predict the SZ power spectrum. The resulting distribution of powers represents the prior probability of a given amount of SZ power, given the current large-angle CMB data, i.e., CMB-normalized predictions for the SZ power spectrum.

3. ANGULAR POWER SPECTRUM OF THE SZ EFFECT

The SZ effect arises from Compton scattering of CMB photons with hot electrons in gas in halos (Sunyaev and Zel'dovich 1980). Many authors have estimated the SZ angular power spectrum C_l using analytic methods or hydrodynamic simulations, with broad agreement between the calculations to within a factor of two at $l < 10^4$ (e.g., see Komatsu and Seljak 2002, for a recent analytic prediction and comparison between analytic methods and numerical simulations; see references therein for previous work). Komatsu and Seljak (2002) find that C_l is sensitive to the baryon density and σ_8 with an approximate scaling of $C_l \propto \sigma_8^7 (\Omega_b h)^2$, but is almost independent of any other cosmological parameters.

We compute C_l as a sum of one-halo Poisson contributions over all halos which can contribute:

$$C_l = g_\nu^2 \int dz \frac{dV}{dz} \int dM \frac{dn(M, z)}{dM} |\tilde{y}_l(M, z)|^2, \quad (1)$$

where $V(z)$ is the comoving volume of the universe at z per steradian, $dn(M, z)/dM$ the comoving dark-matter halo mass function, and g_ν the spectral function of the SZ effect (Sunyaev and Zel'dovich 1980). We ignore the correlated contribution as it is unimportant at $l > 300$ (Komatsu and Kitayama 1999). The physics of gas in halos is encoded in $\tilde{y}_l(M, z)$, the 2D Fourier transform of the Compton y -parameter as a function of halo mass and redshift. The y -parameter is directly proportional to the integrated gas pressure along the line of sight; this leads to $C_l \propto (\Omega_b h)^2$.

We use the method of Komatsu and Seljak (2002) for computing C_l . In brief, we compute a gas-pressure profile using the universal gas-density and temperature profiles derived by Komatsu and Seljak (2001) which make three assumptions about gas in halos: (1) hydrostatic equilibrium between gas pressure and the gravitational force induced by a universal dark-matter density profile (Navarro et al. 1997), (2) the gas density tracing the dark matter density at large radii, as observed in simulated galaxy clusters (e.g., Frenk et al. 1999), and (3) a constant polytropic equation of state for the gas. For the mass function we

use the mass function of Jenkins et al. (2001). This prescription has *no* free parameters, and is in broad agreement with SZ simulations to within a factor of two at $l < 10^4$, agreeing with simulations at least as well as different simulations agree with each other.

Gas cooling, energy feedback, and star formation are not included in our models; Komatsu and Seljak (2002) have argued that these effects are not very important for C_l , which is dominated by gas outside the core of halos. Recently, White et al. (2002) have used hydrodynamic simulations to show that these effects affect C_l by no more than a factor of two. These studies suggest that our theoretical prediction for C_l is accurate to better than a factor of two, and thus MCMC-derived predictions for C_l in the next section can be trusted up to this accuracy.

The SZ fluctuations are highly non-Gaussian, as characterized by large skewness (Seljak et al. 2001; Zhang et al. 2002; White et al. 2002); thus, sampling variance of C_l for the SZ fluctuations differs from that for Gaussian fluctuations, and measurements of C_l in adjacent bins are highly correlated (Cooray 2001). We have to include the non-Gaussianity in the error analysis to interpret observational SZ data correctly. We do this by explicitly including those trispectrum configurations which contribute to the power-spectrum covariance matrix. The error of C_l in a bin of size Δl is given by

$$(\Delta C_l)^2 = f_{\text{sky}}^{-1} \left[\frac{2(C_l + C_l^{\text{N}})^2}{(2l+1)\Delta l} + \frac{T_{ll}}{4\pi} \right], \quad (2)$$

where T_{ll} is the angular trispectrum relevant for the non-Gaussian sampling variance (Cooray 2001),

$$T_{ll} = g_\nu^4 \int dz \frac{dV}{dz} \int dM \frac{dn(M, z)}{dM} |\tilde{y}_l(M, z)|^4. \quad (3)$$

The expression for ΔC_l is correct when both C_l and T_{ll} are sufficiently smooth in l as is the case here. Komatsu and Seljak (2002) have found that the predicted ΔC_l agrees with hydrodynamic simulations to within a factor of two.

For the instrumental-noise power spectrum C_l^{N} we use $l(l+1)C_l^{\text{N}}/(2\pi) = 500, 1000, 1500,$ and $3000 \mu\text{K}^2$ at $l = 1703, 2183, 2630,$ and 3266 for CBI, and 720 and $2000 \mu\text{K}^2$ at $l = 5237$ and 8748 for BIMA. For sky coverage of observations $4\pi f_{\text{sky}}$ we use 1 deg^2 for CBI, and 0.1 deg^2 for BIMA. The bin sizes are $\Delta l = 565, 378, 612, 1000, 2870,$ and 4150 . Although we do not use the exact window functions for the experiments, C_l is very flat at these multipoles, reducing the importance of the window functions.

4. RESULTS

In figure 1 we show the histogram (weighted according to the weights of Lewis and Bridle (2002) for each model) of expected SZ power for CBI and BIMA high- l bins, where the three highest- l CBI bins and the two BIMA bins are averaged to single bins for each experiment. The most likely values for the expected SZ power are less than $100 \mu\text{K}^2$. The weighted mean values are 97 and $86 \mu\text{K}^2$ for CBI and BIMA, respectively, while the median values are 79 and $69 \mu\text{K}^2$. We also find that the analytic approximate scaling of Komatsu and Seljak (2002) at $l \sim 2000 - 6000$, $l(l+1)C_l/(2\pi) \simeq 330 \mu\text{K}^2 \sigma_8^7 (\Omega_b h / 0.035)^2$, gives similar results; the weighted mean is $100 \mu\text{K}^2$, while the median is $88 \mu\text{K}^2$. Bond et al. (2002) find 20–30% lower normalization for the scaling relation at the CBI band, slightly shifting the histogram leftward.




FIG. 1.— Unnormalized probability distribution of the SZ angular power spectrum in CBI (left panel) and BIMA (right panel) high- l averaged windows (solid lines), weighted by the primary CMB likelihood. The gray areas show 95% confidence regions of the CBI (Mason et al. 2002) and BIMA (Dawson et al. 2002) data, assuming Gaussian fluctuations. Note that the true error is larger because of non-Gaussian nature of the SZ fluctuations, and depends on cosmological models. Also shown is the result using analytic approximation of Komatsu and Seljak (2002) for C_l (dashed lines).

For CBI the 95% lower limit (assuming Gaussian fluctuations) to the observed anisotropy is $199 \mu\text{K}^2$, making an SZ interpretation of the CBI measurement only marginally consistent with the CMB-calibrated theories. We show this explicitly in figure 2 where we plot the number of standard deviations of the CBI and BIMA data relative to the predicted levels of SZ fluctuations. We have computed the errors ΔC_l using equation (2). We find that the CBI measurements at $l = 2183$ and 2630 are inconsistent with most of the models that are consistent with the primary CMB data at greater than 2.5 and 2σ levels, respectively. The rest of the CBI bins and the BIMA bins are consistent with the predicted levels of the SZ effect.

The cutoff at the $2.5 - 3\sigma$ level in figure 2 is due to instrumental noise. At the $2.5 - 3\sigma$ level the current data are consistent with no SZ signal. Thus, lower noise experiments could extend the histograms to larger σ values. Note that the largest values of σ in the figure are higher than signal-to-noise ratios of CBI quoted by Mason et al. (2002). This is because ΔC_l depends on the SZ power spectrum; a lower SZ power spectrum leads to lower sample variance, and therefore a smaller ΔC_l . As a result, at the largest σ the instrumental noise determines ΔC_l entirely, making it smaller than the quoted CBI errors.

Which models can account for the data at $l = 2000 - 3000$? We find that those models typically have $\sigma_8 \gtrsim 1$ and $\Omega_b h^2$ higher than the BBN-allowed value of $\Omega_b h^2 = 0.020 \pm 0.002$ (Burles et al. 2001). A simple scaling of $C_l \propto \sigma_8^7 (\Omega_b h)^2$ accounts for this; thus, we need $\sigma_8 \gtrsim 1$ for a given BBN value of $\Omega_b h$ to account for the CBI data by the SZ effect. This conclusion agrees with Bond et al. (2002) and Komatsu and Seljak (2002).

In principle, it would be straightforward to do a joint likelihood analysis of the primary CMB and the SZ effect using CMB data on all angular scales. In practice, the strong non-Gaussianity of the SZ signal, combined with the uncertain field

selection effects (e.g., no bright point sources) make such a procedure still premature. Furthermore, the required $\sigma_8 \gtrsim 1$ is in the tail of the CMB-preferred distribution, so the resolution of the MCMC chain is poor in this region. Importance sampling becomes unreliable in regions where the density of points is low.




FIG. 2.— Number of standard deviations of observed data points relative to predicted SZ fluctuations (each data point corresponds to a panel), weighted by the primary CMB likelihood. The top four panels are the CBI data points (Mason et al. 2002), while the bottom two panels the BIMA data (Dawson et al. 2002). The effects of non-Gaussianity have been included in the error estimation.

5. DISCUSSION

We have presented estimates of the SZ angular power spectrum at $l \gtrsim 2000$ for cosmological models that are consistent with the current CMB anisotropy data at $l \lesssim 2000$. The mode of distribution of the predicted SZ power at $l \sim 3000$ is $50 \mu\text{K}^2$, while we find that the weighted mean is $97 \mu\text{K}^2$ (the median is $78.5 \mu\text{K}^2$), still significantly lower than the reported CBI detection of anisotropy on these angular scales, $508 \mu\text{K}^2$. Moving out from the mode at levels of equal likelihood, the ranges of power enclosed by 68% and 95% of the distribution are $20 - 200 \mu\text{K}^2$ and $5 - 450 \mu\text{K}^2$, respectively. This is effectively a calculation of the primary CMB prior for the SZ power spectrum at the CBI band. When compared to the quoted 68% ($359 - 624 \mu\text{K}^2$) and 95% ($199 - 946 \mu\text{K}^2$) regions of the CBI detection at face value, we find that the CBI detection is only marginally consistent with the SZ effect. The BIMA detection is fully consistent with the SZ effect.

By doing a careful statistical analysis, taking into account the non-Gaussianity of SZ fluctuations and sampling variance, we find that an SZ interpretation of the CBI data is inconsistent at more than $2 - 2.5\sigma$ for those cosmological models which are most consistent with the primary CMB data. The models with the highest SZ power share several characteristics: all models with more than $400 \mu\text{K}^2$ of power but a few exceptions have $0.95 < \sigma_8$ and $0.05 < \Omega_b$. Furthermore, such models prefer a low Hubble constant ($h < 0.7$) and a high matter density ($0.3 < \Omega_{\text{CDM}} + \Omega_b < 0.8$). A reason for a low h is that the primary CMB data tightly constrain $\Omega_b h^2$ while we need a higher $\Omega_b h$ to make the SZ effect larger. We can accomplish this by

increasing Ω_b while reducing h slightly within the HST- h prior. A model with large red tilt ($n_s < 0.95$) can not produce a large amount of SZ power.

The possible presence of radio point sources that partially “fill in” the SZ effect from clusters (Holder 2002) further enhances the possible discrepancy. A reliable determination of σ_8 significantly lower than 1 would be very difficult to reconcile with an SZ interpretation of the measured high- l power.

Uncertainty in estimates for the SZ power spectrum is not yet fully understood, making a more detailed interpretation of the excess power complicated. A major issue is that different simulations have not yet converged, even for adiabatic simulations. Although our current knowledge of missing physics such as gas cooling, star formation, or energy feedback is still limited, these effects appear to not be very important (White et al. 2002) except on very small angular scales ($< 1'$ or $l > 10^4$) where other effects like non-sphericity or merging of halos may also play a role. Nevertheless, the current differences among analytic models, simulations, and estimates of the effects of missing physics are at the level of a factor of two in C_l , while the discrepancy between these predictions and the CBI data is about a factor of five.

Given the strong dependence of C_l on σ_8 , we argue that the discrepancy is due to the difference between a low $\sigma_8 \lesssim 0.8$ favored by the primary CMB and a high $\sigma_8 \gtrsim 1$ favored by the SZ effect (Bond et al. 2002; Komatsu and Seljak 2002). Lahav et al. (2002) and Melchiorri and Silk (2002) have found similarly low σ_8 from the primary CMB data with the same prior on h . *If the excess power is really due to the SZ effect*, then this discrepancy is suggesting that there are some missing components in our analysis. Multi-band SZ observations covering several frequencies will be required to verify the apparent discrepancy.

What is missing in our analysis? The chain that we have used has a strong HST prior on h . Lower values of h reduce the discrepancy, but $h \lesssim 0.4$ would be required to explain the entire difference. Additional components such as massive neutrinos would make the discrepancy worse by driving σ_8 to even lower values. Allowing tensor modes will have competing effects of reducing the overall normalization of scalar modes at large scales but also allowing a blue tilt (higher n_s), leaving the effects on cluster scales largely unchanged. The effect of allowing a general equation of state for the dark energy will be to slightly enhance the SZ fluctuations for a fixed value of σ_8 (Komatsu and Seljak 2002), but to significantly reduce the CMB-preferred value of σ_8 . With the SZ fluctuation power going as

σ_8^7 , the latter effect will dominate and $w > -1$ will generally reduce the expected fluctuation power. The addition of isocurvature fluctuations may help to reconcile the discrepancy, while it significantly expands the allowed range of many cosmological parameters (Trotta et al. 2001).

A very early reionization of the universe ($z_{re} > 20$) will increase the CMB-preferred value of σ_8 , helping to reduce the discrepancy. We find a broad peak in the distribution of models in the chain around $\sigma_8 e^{-\tau} \sim 0.8$, where τ is the Thomson-scattering optical depth of the universe; $\tau \gtrsim 0.22$ or $z_{re} \gtrsim 20$ would comfortably allow $\sigma_8 \gtrsim 1.0$. The discrepancy thus disappears if the universe was reionized early. The current CMB data cannot break the degeneracy between σ_8 and τ , and allow this area of parameter space, although $\tau \gtrsim 0.3$ ($z_{re} \gtrsim 30$) appears to be ruled out. CMB-polarization experiments on large angular scales (e.g., MAP or Planck) should be able to break this degeneracy, and detect the signature of reionization (Zaldarriaga et al. 1997; Eisenstein et al. 1999; Kaplinghat et al. 2002).

We have presented an example of the ease and power of MCMC methods in applying CMB constraints to calculations that include non-trivial dependence on cosmological parameters. It would be worthwhile to investigate the effects of LSS priors, which should make the discrepancy worse by making σ_8 smaller (Lewis and Bridle 2002; Bond et al. 2002). With the strong preference of the high- l measurements for high values of σ_8 and $\Omega_b h$, it would require running a new chain that includes the CBI and BIMA data points, as importance sampling is unreliable in the tails of the current distribution.

Measurements of CMB anisotropy will continue to improve. MCMC methods provide a natural way to incorporate strong constraints on cosmological parameters from CMB experiments into other cosmological studies. Using CMB information to understand the effects of cosmology will allow better understandings of systematic errors in the measurements and insight into important astrophysical processes.

Many of the calculations and some of the text were generously provided by Eiichiro Komatsu, of Princeton University, who could not be a co-author due to considerations of a possible conflict of interest but was an important contributor. We would like to thank David N. Spergel and Uros Seljak for useful discussions. GPH is supported by the W. M. Keck Foundation at the IAS. We are grateful to Sarah Bridle and Antony Lewis for generously providing their MCMC results publicly.

REFERENCES

- Bond, J. R., et al. 2002, ApJ, submitted (astro-ph/0205386)
 Burlles, S., Nollett, K. M., and Turner, M. S. 2001, ApJ, 552, L1
 Christensen, N., Meyer, R., Knox, L., and Luey, B. 2001, Class. Quant. Grav., 18, 2677
 Cole, S., and Kaiser, N. 1988, MNRAS, 233, 637
 Cooray, A. 2001, Phys. Rev. D, 64, 063514
 Dawson, K. S., Holzzapfel, W. L., Carlstrom, J. E., LaRoque, S. J., Miller, A., Nagai, D., and Joy, M. 2002, ApJ, submitted (astro-ph/0206012)
 Eisenstein, D. J., Hu, W., and Tegmark, M. 1999, ApJ, 518, 2–23
 Freedman, W. L., et al. 2001, ApJ, 553, 47
 Frenk, C. S., et al. 1999, ApJ, 525, 554
 Holder, G. P. 2002, ApJ, in press (astro-ph/0205467)
 Hu, W., and Dodelson, S. 2002, ARA&A, in press (astro-ph/0110414)
 Jenkins, A., Frenk, C. S., White, S. D. M., Colberg, J. M., Cole, S., Evrard, A. E., Couchman, H. M. P., and Yoshida, N. MNRAS, 321, 372
 Kaplinghat, M., Chu, M., Haiman, Z., Holder, G. P., Knox, L., and Skordis, C. ApJ, submitted (astro-ph/0207591)
 Komatsu, E., and Kitayama, T. 1999, ApJ, 526, L1
 Komatsu, E., and Seljak, U. 2001, MNRAS, 327, 1353
 Komatsu, E., and Seljak, U. 2002, MNRAS, submitted (astro-ph/0205468)
 Lahav, O., et al. 2002, MNRAS, 333, 961
 Lewis, A., and Bridle, S. 2002, Phys. Rev. D, submitted (astro-ph/0205436)
 Mason, B. S., et al. 2002, ApJ, submitted (astro-ph/0205384)
 Melchiorri, A., and Silk, J. 2002, preprint (astro-ph/0203200)
 Navarro, J. F., Frenk, C. S., and White, S. D. M. 1997, ApJ, 490, 493
 Seljak, U., Burwell, J., and Pen, U. L. 2001, Phys. Rev. D, 63, 063001
 Sunyaev, R. A., and Zel'dovich, Y. B. 1980, ARA&A, 18, 537
 Trotta, R., Riazuelo, A., and Durrer, R. 2001, Phys. Rev. Lett., 87, 231301
 Wang, X., Tegmark, M., and Zaldarriaga, M. 2002, Phys. Rev. D, 65, 123001
 White, M., Hernquist, L., and Springel, V. 2002, ApJ, submitted (astro-ph/0205437)
 Zaldarriaga, M., Spergel, D. N., and Seljak, U. 1997, ApJ, 488, 1
 Zhang, P., Pen, U. L., and Wang, B. 2002, ApJ, submitted (astro-ph/0201375)